

## Semi-empiric Radiative Transfer Modeling of FUSE Stellar Spectra

A. Lobel<sup>1</sup>, E. H. Avrett<sup>1</sup>, and J. P. Aufdenberg<sup>2</sup>

<sup>1</sup>*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street,  
Cambridge 02138 MA, USA*

<sup>2</sup>*National Optical Astronomy Observatory, P.O. Box 26732, Tucson,  
85726 AZ, USA*

**Abstract.** We present an overview of radiative transfer modeling efforts to interpret spectra of a variety of stellar objects observed with *FUSE*. Detailed radiative transfer modeling of high ion emission line profiles of C III and O VI observed in the far-UV spectrum, provides a powerful means to probe the thermal and dynamic properties of high-temperature plasmas in the atmospheres of stars. We model asymmetric emission lines of C III  $\lambda 977$  (and Mg II *h* & *k*) observed in spectra of luminous cool stars such as  $\alpha$  Aqr, to infer the wind- and microturbulence velocity structures of the upper chromosphere. Semi-empiric radiative transfer models that include transition region temperature conditions, are further developed based on detailed fits to O VI resonance emission lines in the supergiant  $\alpha$  Aqr, the classical Cepheid variable  $\beta$  Dor, and to self-absorbed O VI emission lines in the cataclysmic variable SW UMa.

We observe that the C III resonance line profile of  $\alpha$  Aqr assumes a remarkable asymmetric shape, reminiscent of P Cygni type profiles observed in hot luminous supergiants. The model calculations indicate outflow velocities above  $\sim 140$  km s<sup>-1</sup> at kinetic temperatures of 65 kK and higher. Based on detailed model fits to the narrow red-shifted and self-absorbed O VI emission lines of SW UMa we compute that the gas- and electron density exceed the density conditions of the upper solar transition region by about three orders of magnitude. We propose that the large gas density of  $\rho \simeq 1.4 \cdot 10^{-11}$  g cm<sup>-3</sup> favors a region of warm dense plasma of 100 kK  $\leq T_{\text{gas}} \leq 300$  kK that collapses onto the white dwarf with a mass accretion rate of  $1-2 \cdot 10^{15}$  g s<sup>-1</sup> above or between the accretion disk. We discuss how detailed semi-empiric fits to emission lines observed with the high spectral resolution of *FUSE* can provide reliable constraints on the mass loss or mass accretion rates in these objects.

### 1. *FUSE* Observations and Detailed Radiative Transfer Modeling

We present a comparative study of transition region (TR) wind dynamics in  $\alpha$  Aqr (G2 Ib), SW UMa (CV),  $\beta$  Dor (F-G Ia-Iab) and the Sun, to develop semi-empiric radiative transfer models of the thermal structure of the stellar chromosphere and TR, and to determine the velocity and electron density structures. *FUSE* spectra of the hybrid supergiant  $\alpha$  Aqr have been observed for Science Team Program P218 (Dupree et al. 2005). The classical  $\delta$ -Cepheid variable  $\beta$  Dor was observed in Aug. and Oct. 2003 for GI-D107 (PI A. Lobel), while the dwarf nova SW UMa was observed in Nov. 2001 for GI-B074. The left-hand panel of Fig. 1 shows non-LTE radiative transfer fits to C III  $\lambda 977$  through a




Figure 1. Radiative transfer best fit (*solid line*) to C III  $\lambda 977$  in  $\alpha$  Aqr.

semi-empiric model of the chromosphere and lower TR of  $\alpha$  Aqr. The outward decrease of  $\rho$  (*panels right*) produces a scattering core in the computed emission profile (Lobel & Dupree 2000; 2001), which assumes an asymmetric shape due to opacity in a fast accelerating wind. Best multi-level atom model fits are obtained for  $V_{\text{wind}} > 140 \text{ km s}^{-1}$  in the lower TR ( $T \simeq 65 \text{ kK}$ ), which strongly scatters the blue emission line wing. The model signals a supersonic optically thick warm wind in the outer atmosphere of this hybrid supergiant (Lobel & Dupree 2002).

The left-hand panels of Fig. 2 show O VI lines in the cataclysmic variable SW UMa. The lines are remarkably far red-shifted with respect to heliocentric rest. We suggest that the double-peaked narrow profiles result from self-absorbed emission line formation in an infalling region near the white dwarf that is sufficiently optically thick at  $T_{\text{gas}}$  between 100 kK and 300 kK (*panels right*). The depth of the O VI  $\lambda 1032$  central absorption line core can only be computed with electron densities  $N_e$  at least 1100 times larger compared to the solar TR values.




Figure 2. Best fit (*long-dash dotted line*) to O VI  $\lambda 1032$  &  $\lambda 1037$  in SW UMa.




Figure 3. Best fit to solar O VI (*upper left panel*) emission compared to  $\beta$  Dor.

The downflow velocity ( $V_{\text{wind}}$ ) in the model is computed from the conservation of total mass in spherical geometry with  $\dot{M} = 4\pi\rho(r)V_{\text{wind}}(r)(r + R_{\text{WD}})^2$ , using a mass accretion rate  $\dot{M}$  of  $1 \times 10^{15} \text{ g s}^{-1}$ , and  $R_{\text{WD}} = 0.01 R_{\odot}$ . This  $\dot{M}$ -value provides the best fit to the wavelength position of the central self-absorption core where  $V_{\text{wind}} = +70$  to  $+65 \text{ km s}^{-1}$  in the O VI lines formation region.

The upper left-hand panel of Fig. 3 compares the O VI lines we compute with an atmospheric model of the average quiet Sun with SOHO-SUMER observations (*solid dots*). The lines form over a rather small region of  $\sim 10$  model layers with  $100 \text{ kK} \leq T_{\text{gas}} \leq 300 \text{ kK}$ , or  $\sim 200 \text{ km}$  into the solar TR above the upper chromosphere (*panels right*). A best fit to the FWHM and equivalent width of the lines is obtained with projected microturbulence velocities ( $V_{\text{turb}}$ ) increasing from at least  $\sim 30 \text{ km s}^{-1}$  to  $\sim 39 \text{ km s}^{-1}$  over this thermal range (Warren et al. 1997). A best fit to the core position of the O VI  $\lambda 1032$  line is computed with a mean downflow velocity of  $+5 \text{ km s}^{-1}$ . Preliminary atmosphere models for  $\beta$  Dor (*lower left panel*) require  $V_{\text{turb}}$  exceeding  $40 \text{ km s}^{-1}$  in the O VI lines formation region to match the large line widths we also observe in  $\alpha$  Aqr.

**Acknowledgments.** Financial support for this research has been provided by NASA *FUSE* grants GI-D107 and GI-E068.

## References

- Dupree, A. K., Lobel, A., Young, P. R., Ake, T. B., Linsky, J. L., & Redfield S., 2005, ApJ, submitted
- Lobel, A., & Dupree, A. K. 2000, ApJ, 545, 454
- Lobel, A., & Dupree, A. K. 2001, ApJ, 558, 815
- Lobel, A., & Dupree, A. K. 2002, *FUSE* Science and Data Workshop, March 2002, Abstract Poster Talk, p. 60
- Warren, H. P., Mariska, T. J., Wilhelm, K., & Lemaire, P. 1997, ApJ, 484, L91